### § 29.1045

### §29.1045 Order (case).

The state of tobacco with respect to its moisture content.

[42 FR 21092, Apr. 25, 1977. Redesignated at 47 FR 51721, Nov. 17, 1982, 48 FR 29671, June 28, 1983, 49 FR 16756, Apr. 20, 1984, and 51 FR 25027, July 10, 1986]

#### §29.1046 Oxidized (O).

A term applied to tobacco that has deteriorated and turned black during the curing process. Any leaf of which 10 percent or more of its surface has been blackened during the curing process may be described as oxidized. Oxidized tobacco is also known as barn scald or barn rot. (See Rules 23 and 25.)

[42 FR 21092, Apr. 25, 1977. Redesignated at 47 FR 51721, Nov. 17, 1982, 48 FR 29671, June 28, 1983, 49 FR 16756, Apr. 20, 1984, and 51 FR 25027, July 10, 1986]

#### §29.1047 Package.

A hogshead, tierce, case, bale, or other securely enclosed parcel or bundle.

[42 FR 21092, Apr. 25, 1977. Redesignated at 47 FR 51721, Nov. 17, 1982, 48 FR 29671, June 28, 1983, 49 FR 16756, Apr. 20, 1984, and 51 FR 25027, July 10, 1986]

## §29.1048 Packing.

A lot of tobacco consisting of a number of packages submitted as one definite unit for sampling or inspecting. It is represented to contain the same kind of tobacco and has a common identification number or mark on each package.

 $[42\ FR\ 21092,\ Apr.\ 25,\ 1977.\ Redesignated at\ 47\ FR\ 51721,\ Nov.\ 17,\ 1982,\ 48\ FR\ 29671,\ June\ 28,\ 1983,\ 49\ FR\ 16756,\ Apr.\ 20,\ 1984,\ and\ 51\ FR\ 25027,\ July\ 10,\ 1986]$ 

#### §29.1049 Papery.

A term used to describe thin-bodied, oilless tobacco usually associated with whitish-lemon color.

[51 FR 25027, July 10, 1986; 51 FR 28687, Aug. 11, 1986]

# §29.1050 Prematurity.

A condition of growth and development characteristic of the lower leaves of the tobacco plant. Premature leaves have some appearance of ripeness due to a process of starvation caused by translocation of plant food elements from these leaves to other leaves higher on the stalk.

[42 FR 21092, Apr. 25, 1977. Redesignated at 47 FR 51721, Nov. 17, 1982, 48 FR 29671, June 28, 1983, 49 FR 16756, Apr. 20, 1984, and 51 FR 25027, July 10, 1986]

#### §29.1051 Quality.

A division of a group or the second factor of a grade based on the relative degree of one or more elements of quality.

[42 FR 21092, Apr. 25, 1977. Redesignated at 47 FR 51721, Nov. 17, 1982, 48 FR 29671, June 28, 1983, 49 FR 16756, Apr. 20, 1984, and 51 FR 25027, July 10, 1986]

### §29.1052 Raw.

Tobacco as it appears between the time of harvesting and the beginning of the curing process.

[42 FR 21092, Apr. 25, 1977. Redesignated at 47 FR 51721, Nov. 17, 1982, 48 FR 29671, June 28, 1983, 49 FR 16756, Apr. 20, 1984, and 51 FR 25027, July 10, 1986]

#### § 29.1053 Red (R).

A brownish red.

[42 FR 21092, Apr. 25, 1977. Redesignated at 47 FR 51721, Nov. 17, 1982, 48 FR 29671, June 28, 1983, 49 FR 16756, Apr. 20, 1984, and 51 FR 25027. July 10, 1986]

## § 29.1054 Semicured.

Tobacco in the process of being cured or which is partially but not thoroughly cured. Semicured includes tobacco which contains fat stems, swelled stems, frozen tobacco, frozen stems, or stems that have not been thoroughly dried in the curing process. (See Rule 23.)

[42 FR 21092, Apr. 25, 1977. Redesignated at 47 FR 51721, Nov. 17, 1982, 48 FR 29671, June 28, 1983, 49 FR 16756, Apr. 20, 1984, and 51 FR 25027, July 10, 1986]

### § 29.1055 Side.

A certain phase of quality, color, or length as contrasted with some other phase of quality, color, or length; or any peculiar characteristic of tobacco.

 $[42\ FR\ 21092,\ Apr.\ 25,\ 1977.\ Redesignated\ at\ 47\ FR\ 51721,\ Nov.\ 17,\ 1982,\ 48\ FR\ 29671,\ June\ 28,\ 1983,\ 49\ FR\ 16756,\ Apr.\ 20,\ 1984,\ and\ 51\ FR\ 25027,\ July\ 10,\ 1986]$ 

## 88